

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 March 2002 (21.03.2002)

PCT

(10) International Publication Number  
WO 02/22580 A1

(51) International Patent Classification<sup>7</sup>: C07D 211/54,  
211/20, 207/12, 207/08, A61K 31/445, A61P 31/04

(74) Agent: BECKER, Konrad; Novartis AG, Corporate  
Intellectual Property, Patent & Trademark Department,  
CH-4002 Basel (CH).

(21) International Application Number: PCT/EP01/10502

(22) International Filing Date:  
11 September 2001 (11.09.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
0022440.2 13 September 2000 (13.09.2000) GB  
0022439.4 13 September 2000 (13.09.2000) GB  
0024674.4 9 October 2000 (09.10.2000) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,  
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,  
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,  
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,  
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,  
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,  
ZA, ZW.

(71) Applicant (*for all designated States except US*): BIO-  
CHEMIE GESELLSCHAFT M.B.H. [AT/AT]; A-6250  
Kundl (AT).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): ASCHER, Gerd  
[AT/AT]; Daxerfeld 3, A-6250 Kundl (AT). BERNER,  
Heinz [AT/AT]; Geyergasse 2a, A-1180 Wien (AT).  
HILDEBRANDT, Johannes [DE/AT]; St. Laurentgasse  
12, A-2512 Oeynhausen (AT).

Published:

— with international search report

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: ANTIBACTERIALS MUTILINS


(57) Abstract: A compound of formula (I) wherein R and R<sub>2</sub> together with the nitrogen atom to which they are attached form pyrrolidiny or piperidiny, R<sub>1</sub> is a group of formula (II) R<sub>3</sub> and R<sub>3</sub>' are hydrogen, deuterium or halogen, R<sub>4</sub>, R<sub>5</sub> and R<sub>10</sub> are independently of each other hydrogen or alkyl, R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are hydrogen or deuterium; R<sub>9</sub> is amino, alkyl, aryl, heterocyclyl or mercapto; and, if X is oxygen, R<sub>9</sub> is additionally hydrogen; R<sub>10</sub>' is alkyl, X is sulphur, oxygen, NR<sub>10</sub>, or N+(R<sub>10</sub>)<sub>2</sub>; Y is sulphur or oxygen, and m is 0, 1 or 2; with the proviso that, when R and R<sub>2</sub> together with the nitrogen atom to which they are attached form piperidiny, m is 0, Y is S and Y is attached in position 3 of said piperidine ring, that group of formula (I) which is attached to the piperidine ring via the residue Y is either in the (S)-configuration or in the (R)-configuration, preferably in the (S)-configuration which is e.g. useful as antimicrobial, antibacterial.

WO 02/22580 A1

## ANTIBACTERIAL MUTILINS

The present invention relates to compounds having e.g. antimicrobial, such as antibacterial, activity; more specifically the present invention relates to mutilins.

- 5 In one aspect the present invention provides a compound of formula



wherein

R and R<sub>2</sub> together with the nitrogen atom to which they are attached form pyrrolidinyl or piperidinyl,

- 10 R<sub>1</sub> is a group of formula



R<sub>3</sub> and R<sub>3'</sub> are hydrogen, deuterium or halogen,

R<sub>4</sub> is hydrogen or alkyl, e.g. (C<sub>1-4</sub>)alkyl,

R<sub>5</sub> is hydrogen or alkyl, e.g. (C<sub>1-4</sub>)alkyl,

- 15 R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are hydrogen or deuterium;

R<sub>9</sub> is amino, alkyl, aryl, heterocyclyl or mercapto; and, if X is oxygen, R<sub>9</sub> is additionally hydrogen;

R<sub>10</sub> is hydrogen or alkyl, e.g. (C<sub>1-4</sub>)alkyl,

R'<sub>10</sub> is alkyl, e.g. (C<sub>1-4</sub>)alkyl,

- 20 X is sulphur, oxygen, NR<sub>10</sub>, or N<sup>+</sup>(R'<sub>10</sub>)<sub>2</sub> e.g. in the presence of an appropriate anion;

Y is sulphur or oxygen, and

m is 0, 1 or 2;

with the proviso that, when R and R<sub>2</sub> together with the nitrogen atom to which they are attached form piperidinyl, m is 0, Y is S and Y is attached in position 3 of said piperidine ring

that group of formula I which is attached to the piperidine ring via the residue Y is either in the (S)-configuration or in the (R)-configuration, preferably in the (S)-configuration, e.g. with the proviso that, when R and R<sub>2</sub> together with the nitrogen atom to which they are attached form piperidiny, m is 0, Y is S and Y is attached in position 3 of said piperidine ring, a part of R<sub>9</sub> is either in the (S)-configuration or in the (R)-configuration, e.g. if R<sub>9</sub> is alkyl substituted by amino, that amine group is either in the (S)-configuration or in the (R)-configuration; e.g. with the proviso that, if in a compound of formula I m is 0, a part of formula I is either in the (S)-configuration or in the (R)-configuration, e.g. if R<sub>9</sub> is alkyl substituted by amino, that amine group is either in the (S)-configuration or in the (R)-configuration.

An anion in a group of formula N<sup>+</sup>(R'<sub>10</sub>)<sub>2</sub> includes appropriate anions, e.g. anions as conventional in an ammonium group as a counterion.

15 In a compound of formula I  
 R<sub>3</sub>, R'<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are preferably hydrogen;  
 R and R<sub>2</sub> together with the nitrogen atom to which they are attached form pyrrolidine or piperidine (with the proviso as indicated above), e.g. unsubstituted pyrrolidine or piperidine (beside being substituted by a group of formula -C(=X)R<sub>9</sub>), or substituted pyrrolidine or  
 20 piperidine (further substituted beside being substituted by a group of formula -C(=X)R<sub>9</sub>), e.g. substituted by one or more groups which are conventional in organic chemistry, e.g. pleuromutilin chemistry. Preferably pyrrolidine or piperidine are unsubstituted (beside being substituted by a group of formula -C(=X)R<sub>9</sub>)). In a compound of formula I pyrrolidine and piperidine include a group -N(R<sub>1</sub>) and are bound to a group -(CH<sub>2</sub>)<sub>m</sub>-Y. The group -N(R<sub>1</sub>) and  
 25 the group -(CH<sub>2</sub>)<sub>m</sub>-Y may be vicinal or in another position in the pyrrolidine or piperidine ring, e.g. in positions 1,2; 1,3; 1,4; 1,5, and, in case of piperidine, 1,6; and are preferably in positions 1,3 or 1,4 in case of the piperidine ring and in positions 1,2 or 1,3 in case of the pyrrolidine ring;  
 R<sub>9</sub> is preferably alkyl, e.g. (C<sub>1-6</sub>)alkyl, such as (C<sub>1-4</sub>)alkyl, e.g. unsubstituted or substituted  
 30 alkyl, e.g. substituted by groups which are conventional in pleuromutilin chemistry, e.g. one or more amino, heterocyclyl, e.g. including a 5 or 6 membered ring containing 1 or 2 nitrogen atoms; e.g. imidazolyl. If R<sub>9</sub> is alkyl substituted by amino, e.g. and heterocyclyl, R<sub>9</sub> is preferably that part of an amino acid which remains if the carboxylic group is split off, e.g.

the group  $-C(=X)-$  wherein X is oxygen can be regarded as the carbonyl part of said amino acid;

X is preferably oxygen;

Y is preferably sulphur, and

5 m is preferably 0 or 1.

If not otherwise defined herein heterocyclyl includes a 5 or 6 membered ring having 1 to 4 heteroatoms selected from S, O and N; e.g. N; optionally condensed with a further ring (system), e.g. condensed with a phenyl ring; e.g. or condensed with a heterocyclyl ring.

10 Heterocyclyl includes unsubstituted or substituted heterocyclyl, e.g. substituted by groups which are conventional in organic chemistry, e.g. pleuromutilin chemistry. Alkyl includes  $(C_{1-6})$ alkyl, e.g.  $(C_{1-4})$ alkyl. Aryl includes phenyl. Amino includes a free amine group, alkyl- and dialkylamine.

15 In another aspect the present invention provides a compound of formula I, wherein  $R_3, R'_3, R_4, R_5, R_6, R_7$  and  $R_8$  are hydrogen;  $R_1$  is a group of formula




20 R and  $R_2$  together with the nitrogen atom to which they are attached form piperidine or pyrrolidine;  $R_9$  is alkyl; X is oxygen; Y is sulphur; and m is 0 or 1; with the proviso that, when R and  $R_2$  together with the nitrogen atom to which they are attached form piperidiny, m is 0, Y is S and Y is attached in position 3 of said piperidine ring, that group of formula I which is attached to the piperidine ring via the residue Y is either in the (S)-configuration or in the (R)-configuration, preferably in the (S)-configuration;

25 e.g. with the proviso that, when R and  $R_2$  together with the nitrogen atom to which they are attached form piperidiny, m is 0, Y is S and Y is attached in position 3 of said piperidine ring, a part of  $R_9$  is either in the (S)-configuration or in the (R)-configuration, e.g. if  $R_9$  is alkyl substituted by amino, that amine group is either in the (S)-configuration or in the (R)-configuration;


30 e.g. with the proviso that, if in a compound of formula I m is 0, a part of formula I is either in the (S)-configuration or in the (R)-configuration, e.g. if  $R_9$  is alkyl substituted by amino, that amine group is either in the (S)-configuration or in the (R)-configuration.

In another aspect the present invention provides a compound of formula




- wherein  $R_{3p}$ ,  $R'_{3p}$ ,  $R_{6p}$ ,  $R_{7p}$  and  $R_{8p}$  are, index-number correspondingly, as defined above for  $R_3$ ,  $R'_3$ ,  $R_6$ ,  $R_7$  and  $R_8$ ; and  $R_{5p}$  is hydrogen or one or more substituents, preferably hydrogen; and if the group attached to the piperidine ring via the sulphur atom is in position 3 of said piperidine ring and  $R_{5p}$  is hydrogen, then the group attached to the sulphur atom is either in the (S)-configuration or in the (R)-configuration, preferably in the (S)-configuration; e.g. with the proviso that in the group attached to the nitrogen atom of the piperidine ring, the amine group is either in the (S)-configuration or in the (R)-configuration.

In another aspect the present invention provides a compound of formula



- wherein  $R_{3q}$ ,  $R'_{3q}$ ,  $R_{6q}$ ,  $R_{7q}$  and  $R_{8q}$  are, index-number correspondingly, as defined above for  $R_3$ ,  $R'_3$ ,  $R_6$ ,  $R_7$  and  $R_8$ ;  $R_{5q}$  is hydrogen or one or more substituents, preferably hydrogen; and  $R_q$  is that part of an amino acid which remains if the carboxylic group is split off; e.g. including a compound of formula




wherein  $R'_q$  is as defined above for  $R_q$ ,


e.g. with the proviso that in a group  $R_q$  or  $R'_q$ , respectively, the amine group of the amino acid residue is either in the (S)-configuration or in the (R)-configuration.

5

In another aspect the present invention provides a compound of formula



and of formula



10 wherein

$R_{3r}$ ,  $R'_{3r}$ ,  $R_{6r}$ ,  $R_{7r}$  and  $R_{8r}$ , or  $R_{3s}$ ,  $R'_{3s}$ ,  $R_{6s}$ ,  $R_{7s}$  and  $R_{8s}$ , respectively, are, index-number correspondingly, as defined above for  $R_3$ ,  $R'_3$ ,  $R_6$ ,  $R_7$  and  $R_8$ ;

$R_{5r}$ , or  $R_{5s}$ , respectively, is hydrogen or one or more substituents, preferably hydrogen; and

$R_{1r}$ , or  $R_{1s}$ , respectively, is that part of an amino acid which remains if the carboxylic group is  
 15 split off; e.g. wherein in a compound of formula  $I_r$ , the group attached to the piperidine ring

via the sulphur atom is either in the (S)-configuration or in the (R)-configuration; e.g. wherein in a group R<sub>11</sub> or R<sub>18</sub>, respectively, the amine group of the amino acid residue is either in the (S)-configuration or in the (R)-configuration.

- 5 In another aspect the present invention provides the compound, e.g. a compound selected from the group consisting of,
  - 14-O-[(N-(3-Methyl-2-amino-buturyl-piperidin-3(S)-yl)sulfanyl)acetyl]mutilin, e.g. including
  - 14-O-[(N-(3-Methyl-2(R)-amino-buturyl-piperidin-3(S)-yl)sulfanyl)acetyl]mutilin; and
  - 14-O-[(N-(3-Methyl-2(S)-amino-buturyl-piperidin-3(S)-yl)sulfanyl)acetyl]mutilin;
- 10 14-O-[(N-(3-Methyl-2-amino-buturyl-piperidin-4-yl)sulfanyl)acetyl]mutilin, e.g. including
  - 14-O-[(N-(3-Methyl-2(R)-amino-buturyl-piperidin-4-yl)sulfanyl)acetyl]mutilin, and
  - 14-O-[(N-(3-Methyl-2(S)-amino-buturyl-piperidin-4-yl)sulfanyl)acetyl]mutilin;
  - 14-O-[(N-(3-Methyl-2-amino-buturyl)-piperidin-3-yl)-methylsulfanylacetyl]-mutilin, e.g. including
- 15 14-O-[(N-(3-Methyl-2-amino-buturyl)-piperidine-3(S)-yl)-methylsulfanylacetyl]-mutilin, and
  - 14-O-[(N-(3-Methyl-2-amino-buturyl)-piperidine-3(R)-yl)-methylsulfanylacetyl]-mutilin, such as
  - 14-O-[(N-(3-Methyl-2(S)-amino-buturyl)-piperidine-3(S)-yl)-methylsulfanylacetyl]-mutilin, and
  - 14-O-[(N-(3-Methyl-2(R)-amino-buturyl)-piperidine-3(S)-yl)-methylsulfanylacetyl]-mutilin;
  - 14-O-[(N-(3-Methyl-2-amino-buturyl)-pyrrolidine-2-yl)-methylsulfanylacetyl]-mutilin, e.g.
- 20 including
  - 14-O-[(N-(3-Methyl-2-amino-buturyl)-pyrrolidine-2(R)-yl)-methylsulfanylacetyl]-mutilin, and
  - 14-O-[(N-(3-Methyl-2-amino-buturyl)-pyrrolidine-2(S)-yl)-methylsulfanylacetyl]-mutilin, such as
  - 14-O-[(N-(3-Methyl-2(R)-amino-buturyl)-pyrrolidine-2(R)-yl)-methylsulfanylacetyl]-mutilin and
- 25 14-O-[(N-(3-Methyl-2(S)-amino-buturyl)-pyrrolidine-2(R)-yl)-methylsulfanylacetyl]-mutilin,
  - 14-O-[(N-(3-Methyl-2-amino-buturyl)-pyrrolidin-3-yl)sulfanylacetyl]mutilin, e.g. including
  - 14-O-[(N-(3-Methyl-2(R)-amino-buturyl)-pyrrolidine-3-yl)-sulfanylacetyl]-mutilin and
  - 14-O-[(N-(3-Methyl-2(S)-amino-buturyl)-pyrrolidine-3-yl)-sulfanylacetyl]-mutilin;
  - and
- 30 14-O-[(N-histidinyl-pyrrolidin-3-yl)sulfanylacetyl]mutilin, e.g. including
  - 4-O-[(N(R)-histidinyl-pyrrolidin-3-yl)sulfanylacetyl]mutilin, and
  - 4-O-[(N(S)-histidinyl-pyrrolidin-3-yl)sulfanylacetyl]mutilin.

e.g. in free form or in the form of a salt, e.g. a salt with hydrochloric acid; such as a hydrochloride.

14-O-[(N-histidinyl-pyrrolidin-3-yl)sulfanylacetyl]mutiflin is 14-O-[(N-(3-(imidazol-4-yl)-2-amino-propionyl-pyrrolidin-3-yl)sulfanylacetyl]mutiflin.

- 5 Compounds provided by the present invention are hereinafter designated as "compound(s) of the present invention". A compound of the present invention includes a compound in any form, e.g. in free form, in the form of a salt, in the form of a solvate and in the form of a salt and a solvate. A compound of formula I includes a compound of formula I<sub>p</sub>, I<sub>q</sub>, I<sub>q'</sub>, I<sub>r</sub> and I<sub>s</sub>.
- 10 In another aspect the present invention provides a compound of formula I in the form of a salt, or in the form of a salt and in the form of a solvate, or in the form of a solvate.

- A salt of a compound of formula I includes a pharmaceutically acceptable salt, e.g. including a metal salt or an acid addition salt. Metal salts include for example alkali or earth alkali
- 15 salts; acid addition salts include salts of a compound of formula I with an acid, e.g. hydrogen fumaric acid, fumaric acid, naphthalin-1,5-sulphonic acid, hydrochloric acid, deuteriochloric acid; preferably hydrochloric acid or deuteriochloric acid.

- A compound of formula I in free form may be converted into a corresponding compound in the form of a salt; and vice versa. A compound of formula I in free form or in the form of a
- 20 salt and in the form of a solvate may be converted into a corresponding compound in free form or in the form of a salt in unsolvated form; and vice versa.

- A compound of of the present invention may exist in the form of isomers and isomeric mixtures thereof, e.g. optical isomers, cis trans configured isomers. A compound of the
- 25 present invention may e.g. contain asymmetric carbon atoms and may thus exist in the form of diastereoisomeres and mixtures thereof, e.g. epimers. For example in a compound of formula I, that group bound via group -(CH<sub>2</sub>)<sub>m</sub>-Y to the pyrrolidine or piperidine ring may be in the (R)- and in the (S)-configuration, e.g. including mixtures thereof. E.g. if R<sub>9</sub> is alkyl substituted by amine, e.g. R<sub>9</sub> is that part of an amino acid which remains if the carboxylic
- 30 group is split off, said amine group in R<sub>9</sub> may be in the (S)- or in the (R)-configuration, e.g. including mixtures thereof.

Isomeric, diastereoisomeric and epimeric mixtures may be separated as appropriate, e.g. according to a method as conventional, to obtain pure isomers. Pure isomers may also be



produced as appropriate, e.g. according, e.g. analogously, to a method as conventional, e.g. or as described herein.

The present invention includes a compound of the present invention in any isomeric form and in any isomeric mixture, with the proviso that, when R and R<sub>2</sub> together with the nitrogen atom to which they are attached form piperidiny, m is 0, Y is S and Y is attached in position 3 of said piperidine ring, that group of formula I which is attached to the piperidine ring via the residue Y is either in the (S)-configuration or in the (R)-configuration, preferably in the (S)-configuration;

e.g. with the proviso that, when R and R<sub>2</sub> together with the nitrogen atom to which they are attached form piperidiny, m is 0, Y is S and Y is attached in position 3 of said piperidine ring, a part of R<sub>9</sub> is either in the (S)-configuration or in the (R)-configuration, e.g. if R<sub>9</sub> is alkyl substituted by amino, that amine group is either in the (S)-configuration or in the (R)-configuration;


e.g. with the proviso that, if in a compound of formula I m is 0, a part of formula I is either in the (S)-configuration or in the (R)-configuration, e.g. if R<sub>9</sub> is alkyl substituted by amino, that amine group is either in the (S)-configuration or in the (R)-configuration.

Preferably the configuration in the mutilin ring of a compound of formula I is the same as in a naturally produced mutilin.


A compound of the present invention may be obtained as appropriate, e.g. according, e.g. analogously, to a method as conventional, or, as described herein.

In another aspect the present invention provides a process for the production of a compound of formula I as defined above comprising the steps

a. reacting a compound of formula




wherein  $R_3$ ,  $R'_3$ ,  $R_4$  and  $R_5$  are as defined above and  $R_6$ ,  $R_7$  and  $R_8$  are hydrogen, with urea or thiourea and subsequent reduction to obtain a compound of formula



5 wherein Y is as defined above;  $R_3$ ,  $R'_3$ ,  $R_4$  and  $R_5$  are as defined above and  $R_6$ ,  $R_7$  and  $R_8$  are hydrogen,

- b. reacting a compound of formula III as defined in step a. with optionally substituted pyrrolidine, methyl or ethyl pyrrolidine, piperidine, methyl or ethyl piperidine (= methyl-, ethyl- pyrrolidine or piperidine), respectively, carrying at the nitrogen atom a group of formula  $-C(=X)R_9$ , wherein X and  $R_9$  are as defined above, in the form of a reactive derivative, e.g. in the form of a mesylate or a tosylate; to obtain a compound of formula
- 10



which is a compound of formula I wherein  $R$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R'_3$ ,  $R_4$ ,  $R_5$ , Y and m are as defined above and  $R_6$ ,  $R_7$  and  $R_8$  are hydrogen; and, if desired,

- c. introducing deuterium into a compound of formula IV as defined in step b, to obtain a compound of formula I, wherein  $R$ ,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R'_3$ ,  $R_4$ ,  $R_5$ , Y and m are as defined above and  $R_6$ ,  $R_7$  and  $R_8$  are deuterium.
- 15

Groups may be unprotected or protected and may be deprotected in any step, if desired, e.g. according, e.g. analogously to a method as conventional.

E.g. in an optionally substituted pyrrolidine, methyl or ethyl pyrrolidine, piperidine, methyl or ethyl piperidine (= methyl-, ethyl- pyrrolidine or piperidine), respectively, carrying at the nitrogen atom a group of formula  $-C(=X)R_9$ , wherein  $R_9$  is alkyl substituted by amine, said amine group may be protected or unprotected. Appropriate protection groups include e.g.

5 protection groups as conventional, such as tert.butoxycarbonyl (BOC).


If in step b. as defined above the methane- or toluenesulphonic acid residue, respectively, attached to the methyl-, ethyl- pyrrolidine or piperidine ring is in the (R)-configuration a compound of formula IV obtained may be in a form, wherein the group attached to the methyl-, ethyl- pyrrolidine or piperidine ring via the sulphur atom is in the (S)-configuration; If

10 in step b. as defined above the methane- or toluenesulphonic acid residue, respectively, attached to the methyl-, ethyl- pyrrolidine or piperidine ring is in the (S)-configuration a compound of formula IV obtained may be in a form, wherein the group attached to the methyl-, ethyl- pyrrolidine or piperidine ring via the sulphur atom is in the (R)-configuration (Walden inversion).

15 If in step b. as defined above in a group of formula  $-C(=X)R_9$  attached to the methyl-, ethyl- pyrrolidine or piperidine ring used for reaction with a compound of formula III  $R_9$  is alkyl substituted by amine, e.g. that part of an amino acid which remains if the carboxylic group is split off, wherein said amine is in the (R)- configuration, a compound of formula IV is obtained, wherein said amine is in the (R)-configuration. If in step b. as defined above in a


20 group of formula  $-C(=X)R_9$  attached to the methyl-, ethyl- pyrrolidine or piperidine ring used for reaction with a compound of formula III  $R_9$  is alkyl substituted by amine, e.g. that part of an amino acid which remains if the carboxylic group is split off, wherein said amine is in the (S)-configuration, a compound of formula IV is obtained, wherein said amine is in the (S)-configuration.

25 The production of a compound of formula I, wherein  $R_3$  and  $R'_3$  are deuterium or halogen may be e.g. carried out via treatment of a compound of formula



- wherein the carbon atoms carrying  $R_3$  and  $R'_3$ , which both are hydrogen, together form a double bond with deuterium or halogen, e.g.  $F_2$ ,  $Cl_2$ ,  $Br_2$ , to obtain a compound of formula V, wherein  $R_3$  and  $R'_3$  are deuterium or halogen; and further reacting a compound of formula V as appropriate to obtain a compound of formula I wherein  $R_3$  and  $R'_3$  are deuterium or
- 5 halogen.

Preferably a compound of formula II may be obtained from a compound of formula V by reacting a compound of formula V with a compound of formula



- wherein Y,  $R_4$  and  $R_5$  are as defined above and Hal is halogen, preferably bromo, chloro.
- 10 Introduction of deuterium in a compound of formula I may be carried out by treatment of a compound of formula I, wherein  $R_6$ ,  $R_7$  and  $R_8$  are hydrogen, e.g. in the form of a hydrochloride, with deuteriochloric acid (DCI) in appropriate solvent (system); and isolation of a compound of formula I wherein  $R_6$ ,  $R_7$  and  $R_8$  are deuterium. If a compound of formula I is in the form of an acid addition salt, such as a hydrochloride, treatment of said
- 15 salt with DCI may also result in the replacement of the hydrogen of said acid, e.g. in a compound of formula I in the form of a deuteriochloride.

- Any compound described herein, e.g. compounds of formula II, III, IV, V, VI and optionally substituted pyrrolidines, methyl or ethyl pyrrolidines, piperidines, methyl or ethyl piperidines, respectively, are known or may be obtained according, e.g. analogously, to a method as
- 20 conventional, e.g. or as described herein.

- The compounds provided by the present invention including compounds of formula I, hereinafter designated as "active compound(s) of the present invention" exhibit pharmacological activity and are therefore useful as pharmaceuticals.
- 25 For example, the active compounds of the present invention, e.g. compounds of the examples, show antimicrobial, e.g. antibacterial, activity against gram positive bacteria, such as Staphylococci, e.g. Staphylococcus aureus, Streptococci, e.g. Streptococcus pyogenes, Streptococcus pneumoniae, Enterococci, e.g. Enterococcus faecium, as well as against Mycoplasmas, Chlamydia and obligatory anaerobes, e.g. Bacteroides fragilis; in vitro
- 30 In the Agar Dilution Test or Microdilution Test according to National Committee for Clinical Laboratory Standards (NCCLS) 1997, Document M7-A4 Vol.17, No. 2: "Methods for dilution

Antimicrobial Susceptibility Tests for Bacteria that Grow Aerobically – Fourth Edition, Approved Standard"; and in the Anaerobic Bacteria TEST according to National Committee for Clinical Laboratory Standards (NCCLS) VOL. 13, No. 26, M11-A4, Methods for Antimicrobial Susceptibility Testing of Anaerobic Bacteria; Approved Standard; Fourth Edition (1997).

The active compounds of the present invention show antibacterial activity in vitro (MIC ( $\mu\text{g/ml}$ )) in the Agar Dilution Test or in the Microdilution Test from about  $\leq 0.01 \mu\text{g/ml}$  to  $25 \mu\text{g/ml}$ , e.g. against above mentioned bacterial species; and are active against Mycoplasmas and Chlamydia. MIC = minimal inhibitory concentration.

The active compounds of the present invention show e.g. activity in systemic infections of mice determined according to the Handbook of animal models of infection. Edition Oto Zak and Merle A. Sande, Academic Press, 1999), e.g. against Staphylococcus, e.g. when administered parenterally or orally, e.g. at dosages from about 5 to  $150 \text{ mg/kg}$  body weight; e.g. the  $\text{ED}_{50}$  values for the compounds 14-O-[(N-(R)-histidinyl-pyrrolidin-3-yl)sulfanylacetyl]mutilin or 14-O-[(N-(3-methyl-2(R)-amino-butyryl)-pyrrolidine-3-yl)-sulfanylacetyl]mutilin is of about  $11.0 \text{ mg/kg}$  body weight; and the  $\text{ED}_{50}$  values for the compound 14-O-[(N-(3-methyl-2(R)-amino-butyryl)-piperidin-3(S)-yl)sulfanyl]acetyl]mutilin, is in the range of  $7.0 \text{ mg/kg}$  body weight.

$\text{ED}_{50}$  = Effective dosage in  $\text{mg/kg}$  body weight per application by which 50% of the treated animals are protected from death; calculated by Probit analysis ( $n=8$  animals/group).

In another aspect the present invention provides a compound of the present invention for use as a pharmaceutical, preferably as an antimicrobial, such as an antibiotic, e.g. and as an anti-anaerobic.

25

In another aspect the present invention provides a compound of the present invention for use in the preparation of a medicament for the treatment of microbial diseases, for example of diseases caused by bacteria, e.g. selected from Staphylococci, Streptococci, Enterococci; e.g. and of diseases caused by Mycoplasmas, Chlamydia and obligatory anaerobes.

30

Surprisingly the active compounds of the present invention show also activity against strains which are resistant against erythromycin(s), tetracycline(s), e.g. strains including penicillin or multidrug-resistant strains, e.g. of Staphylococcus aureus (MRSA).

In another aspect the present invention provides a method of prophylaxis and treatment of microbial diseases which comprises administering to a subject in need of such treatment an effective amount of a compound of the present invention, e.g. in the form of a pharmaceutical composition.

For prophylaxis and treatment of microbial diseases, the appropriate dosage will, of course, vary depending upon, for example, the active compound of the present invention employed, the host, the mode of administration and the nature and severity of the conditions being treated. However, in general, for satisfactory results in larger mammals, for example humans, an indicated daily dosage is in the range from about 0.5 to 3 g, of an active compound of the present invention conveniently administered, for example, in divided doses up to four times a day.

An active compound of the present invention may be administered by any conventional route, preferably orally, e.g. in form of tablets, powders, capsules, suspensions; e.g. including non-resorbable oral formulations; or parenterally, e.g. in the form of injectable solutions or suspensions; or topically, e.g. in the form of nasal sprays, body solutions, creams, eye drops.

The active compounds of the present invention may be administered in analogous manner, e.g. in similar doses and for similar indications, as erythromycin(s), tetracycline(s).

The active compounds of the present invention may be administered in the form of a pharmaceutically acceptable salt, e.g. an acid addition salt or metal salt; or in free form; optionally in the form of a solvate. The active compounds of the present invention in the form of a salt exhibit the same order of activity as the active compounds of the present invention in free form.

In another aspect the present invention provides a pharmaceutical composition comprising a compound of the present invention in free form or in the form of a pharmaceutically acceptable salt; e.g. and/or in the form of a solvate; in association with at least one pharmaceutical carrier or diluent.

Such compositions may be manufactured according, e.g. analogously, to a method as conventional. Unit dosage form may contain, for example, about 100 mg to about 1 g.

The active compounds of the present invention are additionally suitable as veterinary agents, e.g. veterinary active compounds, e.g. in the prophylaxis and in the treatment of microbial, e.g. bacterial diseases, in animals, such as fowl, pigs and calves; e.g. and for diluting fluids for artificial insemination and for egg-dipping techniques.

5

In another aspect the present invention provides a compound of the present invention for use as a veterinary agent.

10

In another aspect the present invention provides a compound of the present invention for the preparation of a veterinary composition which is useful as a veterinary agent e.g. in the prophylaxis and in the treatment of microbial, e.g. bacterial diseases, in animals.

15

The present invention further provides a veterinary method for the prophylaxis and the treatment of microbial, e.g. bacterial, diseases which comprises administering to a subject in need of such treatment an effective amount of a compound of the present invention, e.g. in the form of a veterinary composition.

20

For use of the active compounds of the present invention as a veterinary agent, the dosage will of course vary depending upon the size and age of the animal and the effect desired; for example for prophylactic treatment relatively low doses would be administered over a longer time period, e.g. 1 to 3 weeks. Preferred doses in drinking water are from 0.0125 to 0.05 weight by volume, particularly 0.0125 to 0.025; and in foodstuffs from 20 to 400 g/metric ton, preferably 20 to 200 g/metric ton.

25

It is preferred to administer the active compounds of the present invention as a veterinary agent to hens in drinking water, to pigs in foodstuff and to calves orally or parenterally, e.g. in the form of oral or parenteral preparations.

30

In the following examples which illustrate the invention temperatures are in degree Celsius and are uncorrected.


The following abbreviations are used:

BOC: tert.butoxycarbonyl

TFA: trifluoroacetic acid

DCC: dicyclohexylcarbodiimide

The numbering of the mutilin cyclus referred to in the examples is given in the following formula:





## Examples

## Example 1

## 14-O-[(N-(3-Methyl-2-amino-buturyl-piperidin-3-yl)sulfanyl)acetyl]mutillin

5 A) 14-O-[(Carbamidoysulfanyl)acetyl]mutillin-tosylate

A solution of 15.2 g of thiourea and 106.4 g of pleuromutillin-22-O-tosylate in 250 ml of acetone is heated under reflux, solvent is removed under reduced pressure and 100 ml of hexane are added. A precipitate forms, is filtrated off and dried.

14-O-[(Carbamidoysulfanyl)acetyl]mutillin-tosylate is obtained.

10 B) 14-Mercapto-acetyl-mutillin

A solution of 4.7 g of sodium pyrosulfite ( $\text{Na}_2\text{S}_2\text{O}_5$ ) in 25 ml of  $\text{H}_2\text{O}$  is added to a solution of 12.2 g of 14-O-[(carbamimidoylsulfanyl)acetyl]mutillin-tosylate in a mixture of 20 ml of ethanol and 35 ml of  $\text{H}_2\text{O}$  (warmed to ca.  $90^\circ$ ). 100 ml of  $\text{CCl}_4$  are added to the reaction mixture obtained and the mixture is heated under reflux. The two-phase system obtained is separated, the organic phase is dried and the solvent is evaporated off.

14-Mercapto-acetyl-mutillin is obtained.

C) N-BOC-3(R)-Hydroxy-piperidine

A suspension of 3.48 g of 3-(R)-hydroxypiperidine, 8.72 g of di-tert.butyl-dicarbonat and 4.0 g of N-metyl-morpholine in 70 ml of dioxane is stirred at room temperature. From the mixture obtained the solvent is evaporated off and the evaporation residue is dissolved in  $\text{CH}_2\text{Cl}_2$  and extracted with 1N HCl. The organic phase is dried and the solvent is evaporated off.

5.08 g of N-BOC-3(R)-Hydroxy-piperidine are obtained which can be used without further puification for further reaction.

25 D) N-BOC-3(R)-methylsulfonyloxy-piperidine

A solution of 5.08 g of N-BOC-3(R)-Hydroxy-piperidine and 8.7 g of methanesulfonic acid anhydride in 100 ml of pyridine is stirred at room temperature. Pyridine is distilled off under high vacuum and the distillation residue obtained is dissolved in  $\text{CH}_2\text{Cl}_2$ , which is extracted with 1N HCl. The organic phase obtained is dried and the solvent is evaporated off to dryness. The residue is purified by chromatography.

3.8 g of N-BOC-3(R)-methylsulfonyloxy-piperidine are obtained.

$^1\text{HNMR}(\text{CDCl}_3)$ : 4.7(m,1H, $\text{CHOSO}_2\text{CH}_3$ ), 3.2-3.6(m,4H,CHN), 3.0(s,3H, $\text{CH}_3\text{SO}_2$ ), 1.4(m,9H,tert.butyl).

**E) 14-O-[(N-BOC-Piperidin-3(S)-yl)-sulfanylacetyl]-mutillin**

- A solution of 1.97 g of 22-mercapto-acetyl-pleuromutillin, 1.39 g of N-BOC-3(R)-methyl-sulfonyloxy-piperidine and 0.12 g of sodium in 50 ml of EtOH is heated to 90°. The solvent of the mixture obtained is evaporated off to dryness under vacuum and the residue is
- 5 subjected to chromatography. 2.5 g of 14-O-[(N-BOC-Piperidin-3(S)-yl)-sulfanylacetyl]-mutillin are obtained.

<sup>1</sup>HNMR (CDCl<sub>3</sub>): 6.45, 5.35, 5,2 (3xm, H<sub>19</sub>, H<sub>20</sub>, H<sub>21</sub>), 5.74 (d, 1H, 5,2Hz, H<sub>14</sub>), 3.35 (d, 1H, H<sub>11</sub>, J=5.2Hz), AB-system: 3.12, 3.18, (J=14.7Hz, H<sub>22</sub>), 3.2, 2.95, 2.65, 2.6 (4xm, CH<sub>2</sub>NCH<sub>2</sub>), 2.85 (m, 1H, SCH), 1.18, 1.45 (2xs, (CH<sub>3</sub>)<sub>15</sub>, (CH<sub>3</sub>)<sub>18</sub>), 0.75, 0.88 (2xd, (CH<sub>3</sub>)<sub>16</sub>, (CH<sub>3</sub>)<sub>17</sub>, J=5.4Hz)

10 **F) 14-O-[(N-(3-Methyl-2(R)-amino-buturyl)-piperidin-3(S)-yl)sulfanyl]acetyl]mutillin in the form of a hydrochloride**

- A solution of 280 mg of 14-O-[(N-BOC-Piperidin-3(S)-yl)-sulfanylacetyl]-mutillin in 20ml methylenchloride and 1ml TFA is stirred at room temperature for ca. 30 minutes and the solvent of the mixture obtained is evaporated off to dryness. To the evaporation residue
- 15 obtained, dissolved in 40 ml of CH<sub>2</sub>Cl<sub>2</sub>, 55 mg of N-metylmorpholine, 110 mg of N-BOC-(R)-valine and 105 mg of DCC are added. The mixture obtained is stirred and precipitated dicyclohexylurea is filtrated off. The solvent from the filtrate obtained is evaporated off and the evaporation residue is subjected to chromatography. 14-O-[(N-(3-Methyl-2(R)-(N-BOC-amino)-buturyl)-piperidin-3(S)-yl)sulfanyl]acetyl]mutillin is obtained and is treated with TFA.
- 20 The solvent of the mixture obtained is evaporated off to dryness and the evaporation residue is treated with etheric HCl. Solid, amorphous 14-O-[(N-(3-methyl-2(R)-amino-buturyl)-piperidin-3(S)-yl)sulfanyl]acetyl]mutillin in the form of a hydrochloride is obtained.
- <sup>1</sup>HMR(d<sub>6</sub>-DMSO, 330K): 6.45, 5.35, 5,2 (3xm, H<sub>19</sub>, H<sub>20</sub>, H<sub>21</sub>), 5.74 (d, 1H, 5,2Hz, H<sub>14</sub>), 5.45 (d, 1H, NH, J=7.8Hz), 4.1 (m, 1H, NHCHCO), 3.35 (d, 1H, H<sub>11</sub>, J=5.2Hz), AB-system: 3.12, 3.18, (J=14.7Hz, H<sub>22</sub>), 3.2, 2.95, 2.65, 2.6 (4xm, CH<sub>2</sub>NCH<sub>2</sub>), 2.8 (m, 1H, SCH), 1.18, 1.45
- 25 (2xs, (CH<sub>3</sub>)<sub>15</sub>, (CH<sub>3</sub>)<sub>18</sub>), 0.75, 0.88 (2xd, (CH<sub>3</sub>)<sub>16</sub>, (CH<sub>3</sub>)<sub>17</sub>, J=5.4Hz), 0.78, 0.84 (2xd, (CH<sub>3</sub>)<sub>2</sub>CH J=6.8Hz)

**Example 2**

30 **14-O-[(N-(3-Methyl-2-amino-buturyl)-piperidin-4-yl)sulfanyl]acetyl]mutillin**

According to the method described in Example 1 but using the appropriate starting materials (e.g. 4-hydroxymethyl-piperidine instead of 3-hydroxymethyl-piperidine), 4-O-[(N-(3-Methyl-2(R)-amino-buturyl)-piperidin-4-yl)sulfanyl]acetyl]mutillin in the form of a hydrochloride is obtained.

<sup>1</sup>HMR(d<sub>6</sub>-DMSO): 8.1(b,3H,NH3+), 6.2-6.4(m,1H,H19), 5.55(d,1H,H14), 5.1-5.2(m,2H,H20), 4.25(m,1H, NCHCO), 4.1,4.25(m,1H,NCH), 3.8-3.95(m,1H,NCH), 3.4(d,1H,H11), 3.0-3.2 (m,2H,NCH), 2.8-2.95(m,1H,sCH), 2.4(m1H,H4), 3.25-3.4(m,2H,SCH2CO), 1.08, 1.39(2xs, (CH3)15,(CH3)18), 0.93,0.98(2xd,6H,(CH3)2CH),0.65,0.85(2xd,6H,(CH3)16,(CH3)17).

5

### Example 3

#### 14-O-[(N-(3-Methyl-2-amino-butyryl)-piperidin-3-yl)-methylsulfanylacetyl]-mutillin

##### A) N-BOC-3(R)-Hydroxymethyl-piperidine

A suspension of 3.48 g of 3-(R)-hydroxymethyl-piperidine, 8.72 g of di-tert.butyl-dicarbonat  
10 and 4.0 g of N-methyl-morpholine in 70 ml of dioxane is stirred for ca. 18 hours at room temperature. From the mixture obtained the solvent is evaporated off and the evaporation residue is dissolved in CH<sub>2</sub>Cl<sub>2</sub> and extracted with 1N HCl. The organic phase is dried and the solvent is evaporated off. 5.08 g of N-BOC-3(R)-hydroxymethyl-piperidine are obtained which may be used for further reaction without further purification.

##### 15 B) N-BOC-3(R)-Methylsulfonyloxymethyl-piperidine

A solution of 5.08 g of N-BOC-3(R)-hydroxymethyl-piperidine and 8.7 g of methanesulfonic acid anhydride in 100 ml of pyridine is stirred at room temperature for ca. 22 hours. From the mixture obtained pyridine is evaporated off (high vacuum) and the evaporation residue obtained is dissolved in CH<sub>2</sub>Cl<sub>2</sub>. The organic phase obtained is extracted with 1N HCl, dried  
20 and the solvent is evaporated off. The evaporation residue is subjected to chromatography. 3.8 g of N-BOC-3(R)-methylsulfonyloxymethyl-piperidine are obtained.

##### C) 14-O-[(N-BOC-Piperidin-3(S)-yl)-methylsulfanylacetyl]-mutillin

A solution of 1.97 g of 22-mercapto-pleuromutillin, 1.39 g of N-BOC-3(R)-methylsulfonyloxymethyl-piperidine and 0.12 g of sodium in 50 ml of EtOH is heated at ca.  
25 90°. The solvent of the mixture obtained is evaporated off and the evaporation residue obtained is subjected to chromatography. 2.5 g of 14-O-[(N-BOC-Piperidin-3(S)-yl)-methylsulfanylacetyl]-mutillin are obtained.

##### D) 14-O-[(N-(3-Methyl-2(S)-amino-butyryl)-piperidine-3(S)-yl)-methylsulfanylacetyl]-mutillin in the form of a hydrochloride

30 A solution of 280 mg of 14-O-[(N-BOC-piperidin-3(S)-yl)-methylsulfanylacetyl]-mutillin in 20 ml of CH<sub>2</sub>Cl<sub>2</sub> and 1ml of TFA is stirred at room temperature for ca. 30 minutes and the solvent of the mixture obtained is evaporated off. The evaporation residue obtained is dissolved in 40 ml of CH<sub>2</sub>Cl<sub>2</sub> and to the solution obtained 55 mg of N-methylmorpholine, 110 mg of N-BOC-(S)-valine and 105 mg of DCC are added and the mixture obtained is stirred.

From the mixture obtained a solid (precipitated dicyclohexylurea) is filtrated off and the filtrate obtained is subjected to chromatography. 14-O-[(N-BOC-(3-Methyl-2(S)-amino-butyl)-piperidine-3(S)-yl)-methylsulfanylacetyl]-mutillin is obtained which is treated with TFA and etheric HCl. 14-O-[(N-(3-Methyl-2(S)-amino-butyl)-piperidine-3(S)-yl)-methylsulfanylacetyl]-mutillin in the form of a hydrochloride is obtained.

<sup>1</sup>HNMR(d<sub>6</sub>-DMSO): 6.1-6.2(m,1H,H19), 5.58 (d,1H,H14), 5.5-5.12(m,2H,H20), 4.2 (m,2H,NCHCO,NCH), 3.75 (m,1H,NCH), 3.42(d,1H,H11), 3.28-3.35(m,2H,SCH<sub>2</sub>CO), 3.1(m,2H, SCH<sub>2</sub>), 1.08,1.36(2xs,6H(CH<sub>3</sub>)<sub>15</sub>, (CH<sub>3</sub>)<sub>18</sub>), 0.95,0.98 (2xd,6H,(CH<sub>3</sub>)<sub>2</sub>CH), 0.65,0.83 (2xd,6H(CH<sub>3</sub>)<sub>16</sub>, (CH<sub>3</sub>)<sub>17</sub>).

10

**Example 4****14-O-[(N-(3-methyl-2-amino-butyl)-pyrrolidine-2-yl)-methylsulfanylacetyl]-mutillin**

According to the method described in Example 3 but using the appropriate starting materials (e.g. 2-(S)-hydroxymethyl-pyrrolidine instead of 3-(R)-hydroxymethyl-piperidine and N-BOC-(R)-valine instead of N-BOC-(S)-valine); 14-O-[(N-(3-methyl-2(R)-amino-butyl)-pyrrolidine-2(R)-yl)-methylsulfanylacetyl]-mutillin in the form of a hydrochloride is obtained.

<sup>1</sup>HNMR(d<sub>6</sub>-DMSO): Rotamere, 8.1(b,3H,NH3), 6.1-6.2(m,1H,H19), 5.52 (d,1H,H14), 5.5-5.12(m,2H,H20), 4.15(m,1H,NCHCO), 3.9(m,1H,NCH), 3.6(m,1H,NCH), 3.42(d,1H,H11), 3.28-3.35(m,2H,SCH<sub>2</sub>CO), 2.68,2.85(2xdd,2H,CHCH<sub>2</sub>S), 1.08,1.36(2xs,6H(CH<sub>3</sub>)<sub>15</sub>, (CH<sub>3</sub>)<sub>18</sub>), 0.95,0.98 (2xd,6H,(CH<sub>3</sub>)<sub>2</sub>CH), 0.65,0.83 (2xd,6H(CH<sub>3</sub>)<sub>16</sub>, (CH<sub>3</sub>)<sub>17</sub>).

**Example 5****14-O-[(N-(3-Methyl-2-amino-butyl)-pyrrolidin-3-yl)sulfanylacetyl]mutillin**

According to the method described in Example 1 but using appropriate starting materials, e.g. N-BOC-3(R)-methylsulfonyloxy-pyrrolidine instead of N-BOC-3(R)-methylsulfonyloxy - piperidine, 14-O-[(N-(3-methyl-2(R)-amino-butyl)-pyrrolidine-3-yl)-sulfanylacetyl]-mutillin in the form of a hydrochloride is obtained.

<sup>1</sup>HNMR(d<sub>6</sub>-DMSO,330K): 6.45,5.35,5.2 (3xm,H<sub>19</sub>,H<sub>20</sub>,H<sub>21</sub>), 5.64 (d,1H,5,2Hz,H<sub>14</sub>), 6.3, (b,1H,NH), 3.95 (m,1H,NHCHCO), 3.35 (d,1H,H<sub>11</sub>,J=5.2Hz), AB-system: 3.0,3.1, (J=14.7Hz,H<sub>22</sub>), 3.2, 2.95, 2.65, 2.6 (4xm,CH<sub>2</sub>NCH<sub>2</sub>), 2.8 (m,1H,SCH), 1.18,1.45, (2xs,(CH<sub>3</sub>)<sub>15</sub>, (CH<sub>3</sub>)<sub>18</sub>), 0.75,0.88 (2xd,(CH<sub>3</sub>)<sub>16</sub>, (CH<sub>3</sub>)<sub>17</sub>,J=5.4Hz), 0.86,0.84 (2xd, (CH<sub>3</sub>)<sub>2</sub>CH, J=6.8Hz).

**Example 6****14-O-[(N-histidiny-pyrrolidin-3-yl)sulfanylacetyl]mutillin**


According to the method described in Example 1 but using appropriate starting materials,

- 5 e.g. N,N'-BOC-(R)-histidine instead of N-BOC-(R)-valin, 4-O-[(N-(R)-histidiny-pyrrolidin-3-yl)sulfanylacetyl]mutillin in the form of a hydrochloride is obtained.

- <sup>1</sup>HNMR (d<sub>6</sub>-DMSO, 330K): 7.45,6.73 (2xs,histidine), 6.25,5.1,5.2 (3xm,H<sub>19</sub>,H<sub>20</sub>,H<sub>21</sub>), 5.59 (d,1H,5,2Hz,H<sub>14</sub>), 5.45 (d,1H,NH,J=7.8Hz), 4.5 (d,1H,NHCHCO,J=4.5Hz), 3.6,3.4 (2xm,2H,NCH<sub>2</sub>CH<sub>2</sub>), 3.35 (d,1H,H<sub>11</sub>,J=5.2Hz), 3.4(m,2H,HisCH<sub>2</sub>), ABX-system: 3.12,3.68, J=14.1Hz,6.5Hz,NCH<sub>2</sub>CHS), 3.2,2.95,2.65,2.6 (4xm,CH<sub>2</sub>NCH<sub>2</sub>), 2.7 (m,1H,SCH), 1.18,1.45 (2xs,(CH<sub>3</sub>)<sub>16</sub>,(CH<sub>3</sub>)<sub>18</sub>), 0.75,0.88 (2xd,(CH<sub>3</sub>)<sub>16</sub>,(CH<sub>3</sub>)<sub>17</sub>,J=5.4Hz).
- 10

## Patent claims

1. A compound of formula



5

wherein

R and R<sub>2</sub> together with the nitrogen atom to which they are attached form pyrrolidinyl or piperidinyl,

R<sub>1</sub> is a group of formula



10

R<sub>3</sub> and R'<sub>3</sub> are hydrogen, deuterium or halogen,

R<sub>4</sub> is hydrogen or alkyl,

R<sub>5</sub> is hydrogen or alkyl,

R<sub>6</sub>, R<sub>7</sub> and R<sub>8</sub> are hydrogen or deuterium;

R<sub>9</sub> is amino, alkyl, aryl, heterocyclyl or mercapto; and, if X is oxygen, R<sub>9</sub> is additionally hydrogen;

15

R<sub>10</sub> is hydrogen or alkyl,

R'<sub>10</sub> is alkyl,

X is sulphur, oxygen, NR<sub>10</sub>, or N<sup>+</sup>(R'<sub>10</sub>)<sub>2</sub>,

Y is sulphur or oxygen, and

20

m is 0, 1 or 2;

with the proviso that, when R and R<sub>2</sub> together with the nitrogen atom to which they are attached form piperidinyl, m is 0, Y is S and Y is attached in position 3 of said piperidine ring that group of formula I which is attached to the piperidine ring via the residue Y is either in the (S)-configuration or in the (R)-configuration, preferably in the (S)-configuration.


25

2. A compound of formula I as defined in claim 1, wherein  
 $R_3, R'_3, R_4, R_5, R_6, R_7$  and  $R_8$  are hydrogen;  
 $R_1$  is a group of formula




- 5  $R$  and  $R_2$  together with the nitrogen atom to which they are attached form piperidine or pyrrolidine;  
 $R_9$  is alkyl;  
 $X$  is oxygen;  
 $Y$  is sulphur; and  
 10  $m$  is 0 or 1;  
 with the proviso that, when  $R$  and  $R_2$  together with the nitrogen atom to which they are attached form piperidinyl,  $m$  is 0,  $Y$  is S and  $Y$  is attached in position 3 of said piperidine ring, that group of formula I which is attached to the piperidine ring via the residue  $Y$  is either in the (S)-configuration or in the (R)-configuration, preferably in the  
 15 (S)-configuration.

3. A compound of formula




- 20 wherein  $R_{3p}, R'_{3p}, R_{6p}, R_{7p}$  and  $R_{8p}$  are, index-number correspondingly, as defined in claim 1 for  $R_3, R'_3, R_6, R_7$  and  $R_8$ ; and  $R_{5p}$  is hydrogen or one or more substituents, and if the group attached to the piperidine ring via the sulphur atom is in position 3 of said piperidine ring and  $R_{6p}$  is hydrogen, then the group attached to the sulphur atom is either in the (S)-configuration or in the (R)-configuration.

## 4. A compound of formula



wherein  $R_{3q}$ ,  $R'_{3q}$ ,  $R_{6q}$ ,  $R_{7q}$  and  $R_{8q}$  are, index-number correspondingly, as defined in claim 1 for  $R_3$ ,  $R'_3$ ,  $R_6$ ,  $R_7$  and  $R_8$ ;  $R_{5q}$  is hydrogen or one or more substituents, preferably hydrogen; and  $R_q$  is that part of an amino acid which remains if the carboxylic group is split off.

## 5. A compound of formula



wherein


$R_{3r}$ ,  $R'_{3r}$ ,  $R_{6r}$ ,  $R_{7r}$  and  $R_{8r}$  are, index-number correspondingly, as defined in claim 1 for  $R_3$ ,  $R'_3$ ,  $R_6$ ,  $R_7$  and  $R_8$ ;

$R_{5r}$  is hydrogen or one or more substituents, and

$R_{1r}$  is that part of an amino acid which remains if the carboxylic group is split off.



6. A compound of formula



wherein

$R_{3s}$ ,  $R'_{3s}$ ,  $R_{6s}$ ,  $R_{7s}$  and  $R_{8s}$  are, index-number correspondingly, as defined in claim 1 for  $R_3$ ,  $R'_3$ ,  $R_6$ ,  $R_7$  and  $R_8$ ;

$R_{5s}$  is hydrogen or one or more substituents, and

$R_{1s}$  is that part of an amino acid which remains if the carboxylic group is split off.

7. A compound of any one of claims 1 to 6 in the form of a salt, or in the form of a salt and in the form of a solvate, or in the form of a solvate.

8. A compound of any one of claims 1 to 7 for use as a pharmaceutical.

9. A method of prophylaxis and treatment of microbial diseases which comprises administering to a subject in need of such treatment an effective amount of a compound of any one of claims 1 to 7.

10. A pharmaceutical composition comprising a compound of any one of claims 1 to 7 in free form or in the form of a pharmaceutically acceptable salt in association with at least one pharmaceutical carrier or diluent.

11. A compound of any one of claims 1 to 7 for use as a veterinary agent.

IPC 7 C07D211/54 C07D211/20 C07D207/12 C07D207/9 A61K31/445  
A61P31/04

#### B. FIELDS SEARCHED

IPC 7 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data, CHEM ABS Data

### C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P, X	WO 01 09095 A (ASCHER GERD ;BERNER HEINZ (AT); BIOCHEMIE GMBH (AT); HILDEBRANDT J) 8 February 2001 (2001-02-08) page 1, line 2 - line 3; claim 1	1-11
Y	DE 28 11 314 A (SANDOZ AG) 27 September 1979 (1979-09-27) page 8, line 17 - line 18; claim 1	1-11
Y	US 4 278 674 A (REINSHAGEN HELLMUTH ET AL) 14 July 1981 (1981-07-14) column 8, line 59 -column 9, line 23; claim 1	1-11
Y	US 4 675 330 A (BERNER HEINZ ET AL) 23 June 1987 (1987-06-23) column 2, line 22 - line 41; claim 1	1-11
	-/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

\* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*G\* document member of the same patent family

Date of the actual completion of the international search

**7 December 2001**

Date of mailing of the international search report

17/12/2001

**Name and mailing address of the ISA**  
European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer \_\_\_\_\_

See1mann, I

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, Citation, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 6 020 368 A (HUNT ERIC ET AL) 1 February 2000 (2000-02-01) column 6, line 33 - line 45; claim 1	1-11
Y	WO 00 37074 A (BROOKS GERALD ;HUNT ERIC (GB); SMITHKLINE BEECHAM PLC (GB)) 29 June 2000 (2000-06-29) page 1, line 6 - line 21; claim 1	1-11
Y	WO 00 27790 A (NAYLOR ANTOINETTE ;HOWARD STEVEN (GB); HUNT ERIC (GB); DABBS STEVE) 18 May 2000 (2000-05-18) page 2, line 1 - line 3; claim 1	1-11
Y	WO 99 21855 A (SANDERSON FRANCIS DOMINIC ;DABBS STEPHEN (GB); HUNT ERIC (GB); FRY) 6 May 1999 (1999-05-06) page 1, line 3 - line 5; claim 1	1-11
Y	EGGER H ET AL: "NEW PLEUROMUTILIN DERIVATIVES WITH ENCHANCED ANTIMICROBIAL ACTIVITYII. STRUCTURE-ACTIVITY CORRELATIONS" JOURNAL OF ANTIBIOTICS, JAPAN ANTIBIOTICS RESEARCH ASSOCIATION. TOKYO, JP, vol. 29, no. 9, September 1976 (1976-09), pages 923-927, XP001022880 ISSN: 0021-8820 table 1	1-11

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 01/10502

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
WO 0109095	A	08-02-2001	AU	6832600 A	19-02-2001
			WO	0109095 A1	08-02-2001
DE 2811314	A	27-09-1979	DE	2811314 A1	27-09-1979
US 4278674	A	14-07-1981	CH	572893 A5	27-02-1976
			CH	568272 A5	31-10-1975
			CH	575375 A5	14-05-1976
			CH	560180 A5	27-03-1975
			CH	559719 A5	14-03-1975
			AT	335054 B	25-02-1977
			AT	845872 A	15-06-1976
			BE	789629 A1	03-04-1973
			BG	21386 A3	20-05-1976
			BG	20773 A3	20-12-1975
			CA	979447 A1	09-12-1975
			CS	219306 B2	25-03-1983
			CY	987 A	02-08-1979
			DD	100706 A5	05-10-1973
			DE	2248237 A1	12-04-1973
			DE	2265666 C2	03-02-1983
			DK	139720 B	02-04-1979
			DK	498076 A ,B,	03-11-1976
			DK	498176 A ,B,	03-11-1976
			FI	58635 B	28-11-1980
			FI	794060 A ,B,	27-12-1979
			FI	794061 A ,B,	27-12-1979
			FR	2157828 A1	08-06-1973
			GB	1410506 A	15-10-1975
			GB	1410505 A	15-10-1975
			HK	25379 A	27-04-1979
			HU	168436 B	28-04-1976
			IE	37699 B1	28-09-1977
			IE	37700 B1	28-09-1977
			IL	40494 A	30-01-1976
			IL	47262 A	30-01-1976
			JP	48044244 A	26-06-1973
			JP	57007147 B	09-02-1982
			MY	10379 A	31-12-1979
			NL	7213254 A ,B,	09-04-1973
			NO	136459 B	31-05-1977
			NO	136837 B	08-08-1977
			NO	137440 B	21-11-1977
			OA	4192 A	31-12-1979
			PH	11870 A	04-08-1978
			RO	67478 A1	15-03-1980
			RO	67479 A1	21-06-1981
			SE	398348 B	19-12-1977
			SE	424074 B	28-06-1982
			SE	7509456 A	26-08-1975
			SE	419984 B	07-09-1981
			SE	7509457 A	26-08-1975
			US	4107164 A	15-08-1978
			US	4208326 A	17-06-1980
			YU	249772 A1	27-04-1983
US 4675330	A	23-06-1987	DE	3405632 A1	22-08-1985
			DE	3413708 A1	24-10-1985

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4675330	A	AT 29132 T	15-09-1987
		AU 585209 B2	15-06-1989
		AU 3877385 A	12-09-1985
		CA 1249281 A1	24-01-1989
		CY 1565 A	17-05-1991
		DD 253616 A5	27-01-1988
		DE 3560511 D1	01-10-1987
		DK 72885 A ,B,	18-08-1985
		EP 0153277 A2	28-08-1985
		ES 540464 D0	16-04-1986
		ES 8606262 A1	01-10-1986
		FI 850654 A ,B,	18-08-1985
		FI 921848 A ,B,	24-04-1992
		GR 850415 A1	18-06-1985
		HK 103690 A	21-12-1990
		HU 39155 A2	28-08-1986
		IE 58232 B	11-08-1993
		IL 74355 A	29-04-1988
		JP 1886041 C	22-11-1994
		JP 6008278 B	02-02-1994
		JP 60199870 A	09-10-1985
		KR 9201990 B1	09-03-1992
		LV 5764 A4	20-12-1996
		NZ 211131 A	30-08-1988
		PH 21588 A	11-12-1987
		PL 251972 A1	03-12-1985
		SG 69290 G	23-11-1990
		SU 1582985 A3	30-07-1990
		ZA 8501190 A	24-09-1986
US 6020368	A 01-02-2000	AU 715229 B2	20-01-2000
		AU 1307897 A	01-08-1997
		AU 4203697 A	25-02-1998
		BG 102600 A	30-09-1999
		BR 9612426 A	13-07-1999
		BR 9711008 A	17-08-1999
		CN 1231665 A	13-10-1999
		CZ 9900293 A3	16-06-1999
		WO 9805659 A1	12-02-1998
		EP 0874809 A1	04-11-1998
		EP 0934316 A1	11-08-1999
		HU 0001741 A2	28-10-2000
		JP 2000515532 T	21-11-2000
		JP 2000503642 T	28-03-2000
		NO 983074 A	31-08-1998
		NO 990463 A	01-02-1999
		PL 327737 A1	21-12-1998
		PL 331470 A1	19-07-1999
		SK 91098 A3	11-02-1999
		TR 9900194 T2	22-03-1999
		CA 2240467 A1	17-07-1997
		CN 1214039 A	14-04-1999
		CZ 9802124 A3	16-12-1998
		WO 9725309 A1	17-07-1997
		HU 9900973 A2	30-08-1999
		TR 9801282 T2	21-12-1998
		US 6239175 B1	29-05-2001
		ZA 9700017 A	02-07-1998

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
US 6020368	A		AP 872 A		28-09-2000
			US 6121281 A		19-09-2000
			ZA 9706817 A		01-02-1999
WO 0037074	A	29-06-2000	AU 3036200 A		12-07-2000
			WO 0037074 A1		29-06-2000
WO 0027790	A	18-05-2000	WO 0027790 A1		18-05-2000
WO 9921855	A	06-05-1999	AU 9636198 A		17-05-1999
			BR 9814747 A		20-11-2001
			CN 1283197 T		07-02-2001
			EP 1028961 A1		23-08-2000
			WO 9921855 A1		06-05-1999
			HU 0004040 A2		28-05-2001
			NO 20002173 A		05-06-2000
			PL 340254 A1		29-01-2001
			TR 200001203 T2		21-08-2000
			US 6281226 B1		28-08-2001